



Multi Daily Injection



√ How do insulin pens work?



✓ Action Times for Insulins



✓ Doses:

		Dr.	
Basal	LIO CLI	luo i	
Rayar	1111	1111	
Dasai	11134	1111.	

Works for 24 hours and never stops, it controls sugar before meals, during fasting and sleep

The Basal Insulin like the pancreas



Given every day at the same hour Example: 6 pm every day

Also, remember that it has nothing to do with meals. It is always given even during illness or vomiting.

Dose:		

✓ Doses:

Rapid-acting insulin(Meals):	Rag	oid-a	cting	insulin(Meals):	
------------------------------	-----	-------	-------	----------	-------	----	--


It works from 2 to 8 hours, it has a peak that causes a decrease in sugar. Taken with meals, correction and elevations. Controls the sugar level two hours after the meal.




Dose:
Breakfast
Lunch
Dinner
It is given with the meal and can be given immediately after the meal if the child is not sure that he will
eat or not

✓ Required readings for blood sugar:

These readings are controlled







✓ Correction doses and correction times:

They are extra doses given from Rapid-acting insulin (meals) in the case of a high blood sugar level.

Correction times:

- Before meals
- Two hours after a meal
- Between meals

- ✓ Insulin Sensitivity Factor (ISF): It is the amount of decrease in a glucose reading due to taking a dose equal to one unit of insulin.
- (The device's sugar reading) is subtracted from it

✓ Method of calculating the correction dose:

(the upper limit of the required reading) and the result is divided by (the insulin sensitivity factor).

(the upper limit of the required reading – The device's sugar reading)
	= Correction
the insulin sensitivity	

Notes:

Sources and References:

All icons rights for flaticon

Review and audit:

The content of this booklet has been reviewed by pediatric endocrinology and diabetes consultants at King Fahad University Hospital

Health Awareness Unit

IAU-21-187

